## Rhodora

## JOURNAL OF THE NEW ENGLAND BOTANICAL CLUB

NORTON H. NICKERSON, Editor-in-Chief JOAN Y. NICKERSON, Managing Editor

A. LINN BOGLE WILLIAM D. COUNTRYMAN GARRETT E. CROW RICHARD A. FRALICK GERALD J. GASTONY NORTON G. MILLER

ROBERT T. WILCE

**VOLUME 86** 

1984

The New England Botanical Club, Inc. Harvard University Herbaria, 22 Divinity Ave., Cambridge, Mass. 02138



## **INDEX TO VOLUME 86**

New scientific names are in bold face.

- Adams, Ralph M. see Sauleda, Ruben P. Additions to the flora of Cape Breton Highlands National Park, Nova Scotia 67-71
- Additions to the flora of Ulster County, New York 89-94
- Adirondack mountain summit. A 24year comparison of the vegetation of,
- Allred, Kelly W. Studies in the Aristida (Gramineae) of the southeastern United States I. Spikelet variation in A. purpurescens, A. tenuispica and A. virgata 73-77
- Angelo, Ray and Bruce A. Sorrie. Nantucket field trip of the New England Botanical Club 117
- Annotated list of Minnesota Carices, with phytogeographical and ecological notes 151-231. Floristics 152; annotated list 169; doubtful and excluded taxa 200; distribution maps 203-226
- Aristida (Gramineae) of the southeastern United States I. Spikelet variation in A. purpurescens, A. tenuispica and A. virgata. Studies in the, 73-77
- Aristolochia tomentosa Sims established at two western Massachusetts sites 115
- Arnica (Compositae) subgenus Austromontana. Taxonomy of, 239-309
- Bates, Vernon and Paul Lewis. Rediscovery of Stylisma humistrata (Convolvulaceae) in Tennessee 393-394
- Book review: Where have all the wildflowers gone? 237
- Broughtonia R. Br., Cattleyopsis Lem. and Laeliopsis Lindl. A reappraisal of the orchid genera, 445
- Broughtonia ×jamaicensis Sauleda and Adams, hybr. nov. 453

- Burk, C. John. Aristolochia tomentosa Sims established at two western Massachusetts sites 115
- Burk, C. John. Geranium nepalense var. Thunbergii and G. sibiricum naturalized in western Massachusetts 113
- Burk, C. John see Holland; Marjorie M.
- Campbell, Christopher S. see Famous, Norman C.
- Cape Breton Highlands National Park, Nova Scotia. Additions to the flora of. 67-71
- Catling, Paul M. A re-evaluation of Spiranthes ×steigeri Correll 469
- Cat-tail (Typha latifolia L.) marsh. The effect of power utility right-of-way construction on, 389-391
- Cattleyopsis cubensis (Lindl.) Sauleda & Adams, comb. nov. 455
- Cattleyopsis Lem. and Lailiopsis Lindl.
  A reappraisal of the orchid genera
  Broughtonia R. Br., 445
- Cayman Islands. A new species of Phyllanthus (Euphorbiaceae) from the, 121-125
- Central America. A revision of the genus Kyllinga Rottb. (Cyperaceae) in Mexico and, 507
- Central American species of Cerastium (Caryophyllaceae). A revision of the Mexican and, 339-379
- Cerastium (Caryophyllaceae). A revision of the Mexican and Central American species of, 339-379
- Cerastium cuchumatanense D. A. Good, sp. nov. 349; C. sinaloense D. A. Good, sp. nov. 367; C. tolucense D. A. Good, sp. nov. 371; fig. 350
- Chaptalia (Asteraceae: Mutisieae) from Mexico. A new widespread species of, 127-130

- Chaptalia transiliens Nesom, sp. nov.
- Chromosome number in the genus Stellaria (Caryophyllaceae). A new basic, 417
- Connecticut River oxbow swamp forests.

  The herb strata of three, 397
- Croomia pauciflora (Stemonaceae). Notes on, 131-137
- Denford, Keith E. see Wolf, Steven J. Dentaria laciniata Muhl. (Brassicaceae) reported from Maine. Note on the first
- station of, 233 Distribution and ecological characteristics of Ironwood, Ostrya virginiana (Miller) K. Koch, in northeastern Nova Scotia 139-149
- Distributions. Lomatogonium rotatum (Gentianaceae) and Primula laurentiana (Primulaceae) in Maine: new localities and general, 425
- Domville, Mary. Additions to the flora of Ulster County, New York 89-94
- Ecological notes. Annotated list of Minnesota Carices, with phytogeographical and, 151-231
- cal and, 151-231
  Effect of power utility right-of-way construction on cat-tail (Typha latifolia L.) marsh. The, 389-391
- Environmental corollaries of Panax quinquefolium (Araliaceae) in Delaware County, New York. Population structure and, 431
- Erratum, October 1983 issue (Vol 85, No. 844), Dent and Adams 119
- Famous, Norman C. and Christopher S. Campbell. Lomatogonium rotatum (Gentianaceae) and Primula laurentiana (Primulaceae) in Maine: new localities and general distributions 425
- Farrar, Donald R. see Parks, James C.
- First report of the fern genus Vittaria in New York. A, 421

- Flora boreali-americana of André Michaux. The type localities of the, 1-66
- Florida. A new species of Ziziphus (Rhamnaceae) from, 381-387
- Geranium nepalense var. Thunbergii and G. sibiricum naturalized in western Massachusetts 113
- Geranium sibiricum naturalized in western Massachusetts. Geranium nepalense var. Thunbergii and, 113
- Good, David A. A revision of the Mexican and Central American species of Cerastium (Caryophyllaceae) 339-379
- Gratiola viscidula Pennell (Scrophulariaceae). Infraspecific variation in, 79-87
- Greenidge, K. N. H. Distribution and ecological characteristics of Ironwood, Ostrya virginiana (Miller) K. Koch, in northeastern Nova Scotia 139-149
- Halenia alleniana Standl. ex Wilbur, sp. nov. 326; H. crumiana Wilbur, sp. nov. 321
- Halenia (Gentianaceae) in Mexico. A synopsis of the genus, 311-337
- Hall, David W. see Judd, Walter S.
  Herb strata of three Connecticut River oxbow swamp forests. The, 397. Floristic trends and comparisons 409;
  Study sites: Hatfield oxbow 405, Ned's Ditch 406, Whately oxbow 409; Table: Composition of herb strata: herbs and vines 402, woody seedlings 404
- Hellquist, C. Barre. Observations on Potamogeton hillii Morong in North America 101-111
- Hinds, Harold L. Additions to the flora of Cape Breton Highlands National Park, Nova Scotia 67-71
- Holland, Marjorie M. and C. John Burk. The herb strata of three Connecticut River oxbow swamp forests 397

- Infraspecific variation in Gratiola viscidula Pennell (Scrophulariaceae) 79-87
- Ironwood, Ostrya virginiana (Miller) K. Koch, in northeastern Nova Scotia. Distribution and ecological characteristics of, 139-149
- Judd, Walter S. and David W. Hall. A new species of Ziziphus (Rhamnaceae) from Florida 381-387
- Ketchledge, E. H. and R. E. Leonard. A 24-year comparison of the vegetation of an Adirondack mountain summit 430
- Kyllinga Rottb. (Cyperaceae) in Mexico and Central America. A revision of the genus, 507
- Laeliopsis Lindl. A reappraisal of the orchid genera Broughtonia R. Br., Cattleyopsis Lem, and, 445
- Lassetter, J. Stuart. Taxonomy of the Vicia ludoviciana complex 475
- Leonard, R. E. see Ketchledge, E. H. Lewis, Paul see Bates, Vernon
- Lewis, Walter H. Population structure and environmental corollaries of Panax quinquefolium (Araliaceae) in Delaware County, New York 434
- Lomatogonium rotatum (Gentianaceae) and Primula laurentiana (Primulaceae) in Maine: new localities and general distributions 425
- Maine: new localities and general distributions. Lomatogonium rotatum (Gentianaceae) and Primula laurentiana (Primulaceae) in, 425
- Maine. Note on the first station of Dentaria laciniata Muhl. (Brassicaceae) reported from, 233
- Massachusetts, Geranium nepalense var. Thunbergii and G. sibiricum naturalized in western. 113
- Massachusetts sites. Aristolichia tomentosa Sims established at two western, 115

- Medeolaria farlowii in New England. Two new localities for, 235
- Mexican and Central American species of Cerastium (Caryophyllaceae). A revision of the, 339-379
- Mexico. A new, widespread species of Chaptalia (Asteraceae: Mutisieae) from, 127-130
- Mexico. A synopsis of the genus Halenia (Gentianaceae) in, 311-337
- Mexico and Central America. A revision of the genus Kyllinga Rottb. (Cyperaceae) in, 507
- Michaux, André. The type localities of the Flora boreali-americana of, 1-66
- Minnesota Carices, with phytogeographical and ecological notes. Annotated list of, 151-231
- Morton, J.K. A new basic chromosome number in the genus Stellaria (Caryophyllaceae) 417
- Nantucket field trip of the New England Botanical Club 117
- NEBC Research Award Notice 126, 395, 541
- Nesom, Guy L. A new, widespread species of Chaptalia (Asteraceae: Mutisieae) from Mexico 127-130
- New basic chromosome number in the genus Stellaria (Caryophyllaceae). A, 417
- New England. Two new localities for Medeolaria farlowii in, 235
- New England Botanical Club. Nantucket field trip of the, 117
- New England Notes:
  - Aristolochia tomentosa Sims established at two western Massachusetts sites 115
  - Geranium nepalense var. Thunbergii and G. sibiricum naturalized in western Massachusetts 113
  - Nantucket field trip of the New England Botanical Club 117
  - Note on the first station of Dentaria laciniata Muhl. (Brassicaceae) reported from Maine 233

- The re-discovery of Solidago odora Ait. (Asteraceae) in Vermont 539 Two new localities for Medeolaria farlowii in New England 235
- New localities and general distributions.

  Lomatogonium rotatum (Gentianaceae) and Primula laurentiana (Primulaceae) in Maine: 425
- New species of Phyllanthus (Euphorbiaceae) from the Cayman Islands. A, 121-125
- New species of Ziziphus (Rhamnaceae) from Florida. A, 381-387
- New, widespread species of Chaptalia (Asteraceae: Mutisieae) from Mexico. A, 127-130
- New York. A first report of the fern genus Vittaria in, 421
- New York. Additions to the flora of Ulster County, 89-94
- New York. Population structure and environmental corollaries of Panax quinquefolium (Araliaceae) in Delaware County, 431
- Nickerson, N. H. see Thibodeau, F. R.
- North America. Observations on Potamogeton hillii Morong in. 101-111
- Note on the first station of Dentaria laciniata Muhl. (Brassicaceae) reported from Maine 233
- Notes on Croomia pauciflora (Stemonaceae) 131-137
- Nova Scotia. Additions to the flora of Cape Breton Highlands National Park, 67-71
- Nova Scotia. Distribution and ecological characteristics of Ironwood, Ostrya virginiana (Miller) K. Koch, in northeastern, 139-149
- Observations on Potamogeton hillii Morong in North America 101-111
- Ostrya virginiana (Miller) K. Koch, in northeastern Nova Scotia. Distribution and ecological characteristics of Ironwood. 139-149

- Ownbey, Gerald B. see Wheeler, Gerald A.
- Panax quinquefolium (Araliaceae) in Delaware County, New York. Population structure and environmental corollaries of, 431
- Parks, James C. and Donald R. Farrar. A first report of the fern genus Vittaria in New York 421
- Pfister, Donald H. Two new localities for Medeolaria farlowii in New England 235
- Phyllanthus caymanensis Webster & Proctor, sp. nov. 121; figs. 122, 124
- Phytogeographical and ecological notes. Annotated list of Minnesota Carices, with, 151-231
- Population structure and environmental corollaries of Panax quinquefolium (Araliaceae) in Delaware County, New York 431
- Potamogeton hillii Morong in North America. Observations on, 101-111
- Primula laurentiana (Primulaceae) in Maine: new localities and general distributions. Lomatogonium rotatum (Gentianaceae) and, 425
- Proctor, George R. see Webster, Grady L.
- Reappraisal of the orchid genera Broughtonia R. Br., Cattleyopsis Lem. and Laeliopsis Lindl. A, 445. Key to the genera 447; B. sanguinea 449; negrilensis 451, ×jamaicensis Sauleda & Adams hybr. nov. 453; C. cubensis Sauleda & Adams comb. nov. 455, lindenii 458, ortgiesiana 461; L. domingensis 464
- Rediscovery of Stylisma humistrata (Convolvulaceae) in Tennessee 393-394
- Re-discovery of Solidago odora Ait. (Asteraceae) in Vermont. The, 539
- Re-evaluation of Spiranthes ×steigeri Correll. A, 469

Reinartz, James A. Verbascum densiflorum in southeast Wisconsin 95-99

Revision of the genus Kyllinga Rottb. (Cyperaceae) in Mexico and Central America. A, 507. Key to the species 513; brevifolia 522; nudiceps 530; odorata 518; pumila 514; tibialis 527; vaginata 525

Revision of the Mexican and Central American species of Cerastium (Caryophyllaceae). A, 339–379. C. axillare 346; barberi 346; brachypodum 348; cuchumatanense D. A. Good, sp. nov. 349; glomeratum 352; guatemalense 355; juniperorum 357; madrense 358; nutans 359; orithales 362; purpusii 364; ramigerum 365; sinaloense D. A. Good, sp. nov. 367; sordidum 368; texanum 369; tolucense D. A. Good, sp. nov. 371; triviale 372; vulcanicum 374; excluded species 375; key to the species 341

Right-of-way construction on cat-tail (Typha latifolia L.) marsh. The effect of power utility, 389-391

Rooney, Sally C. Note on the first station of Dentaria laciniata Muhl. (Brassicaceae) reported from Maine 233

Sauleda, Ruben P. and Ralph M. Adams. A reappraisal of the orchid genera Broughtonia R. Br., Cattleyopsis Lem. and Laeliopsis Lindl. 445

Solidago odora Ait. (Asteraceae) in Vermont. The re-discovery of, 539

Sorrie, Bruce A. see Angelo, Ray Spiranthes ×steigeri Correll. A reevaluation of, 469

Spooner, David M. Infraspecific variation in Gratiola viscidula Pennell (Scrophulariaceae) 79-87

Statement of ownership, inside back cover, No. 848

Stellaria (Caryophyllaceae). A new basic chromosome number in the genus, 417

Studies in the Aristida (Gramineae) of the southeastern United States I. Spikelet variation in A. purpurescens, A. tenuispica, and A. virgata 73-77 Stylisma humistrata (Convolvulaceae) in Tennessee. Rediscovery of, 393-394 Swamp forests. The herb strata of three Connecticut River oxbow, 397

Synopsis of the genus Halenia (Gentianaceae) in Mexico. A, 311-337. H. alata 327; alleniana Wilbur sp. nov. 326; brevicornia 315; conzattii 324; crumiana Wilbur sp. nov. 321; decumbens 332; hintonii 331; palmeri 323; plantaginea 334; pringlei 328; recurva 320; schiedeana 325; key to the species 313

Taxonomy of Arnica (Compositae) subgenus Austromontana 239-309. A. cernua 258; cordifolia 262; discoidea 272; gracilis 277; latifolia 282; nevadensis 290; spathulata 295; venosa 300; viscosa 303; chromosome numbers 241; excluded taxa 306; key to the species 257; morphology and taxonomic criteria 242; phylogeny and phytogeography 248

Taxonomy of the Vicia ludoviciana complex (Leguminosae) 475. Key to subspecies and races 486; § leavenworthii 499; § ludoviciana 487

Tennessee. Rediscovery of Stylisma humistrata (Convolvulaceae) in, 393-394 Thibodeau, F. R. and N. H. Nickerson. The effect of power utility right-ofway construction on cat-tail (Typha latifolia L.) marsh 389-391

Tucker, Gordon C. A revision of the genus Kyllinga Rottb. (Cyperaceae) in Mexico and Central America 507

Twenty-four year comparison of the vegetation of an Adirondack mountain summit. A, 439

Two new localities for Medeolaria farlowii in New England 235

Type localities of the Flora borealiamericana of André Michaux 1-66. Michaux Vol. I 6; Michaux Vol. II 40

- Ulster County, New York. Additions to the flora of, 89-94
- United States. Studies in the Aristida (Gramineae) of the southeastern, 73-77
- Uttal, Leonard J. The type localities of the Flora boreali-americana of André Michaux 1-66
- Verbascum densiflorum in southeast Wisconsin 95-99
- Vermont. The re-discovery of Solidago odora Ait. (Asteraceae) in, 539
- Vicia ludoviciana complex (Leguminosae). Taxonomy of the, 475
- Vittaria in New York. A first report of the fern genus, 421
- Webster, Grady L. and George R. Proctor. A new species of Phyllanthus (Euphorbiaceae) from the Cayman Islands 121-125

- Wheeler, Gerald A. and Gerald B. Ownbey. Annotated list of Minnesota Carices, with phytogeographical and ecological notes 151-231
- Whetstone, R. David. Notes on Croomia pauciflora (Stemonaceae) 131-137
- Wilbur, Robert L. A synopsis of the genus Halenia (Gentianaceae) in Mexico. 311-337
- Wisconsin. Verbascum densiflorum in southeast, 95-99
- Wolf, Steven J. and Keith E. Denford. Taxonomy of Arnica (Compositae) subgenus Austromontana. 239-309
- Zika, Peter F. The re-discovery of Solidago odora Ait. (Asteraceae) in Vermont 539
- Ziziphus celata Judd and Hall, sp. nov. 382, fig. 384
- Ziziphus (Rhamnaceae) from Florida. A new species of, 381-387





